

Wikiprint Book

Title: EwEugForagingTimeAndPredationRisk

Subject: Ecopath Developer Site - EwEugForagingTimeAndPredationRisk

Version: 9

Date: 2024-04-26 06:36:28

Table of Contents

3.6 Foraging time and predation risk	3
--------------------------------------	---

3.6 Foraging time and predation risk

The food consumption prediction relationship in Eq. 52 (see [Predicting consumption?](#)) contains two parameters that directly influence the time spent feeding and the predation risk that feeding may entail: a_{ij} and v_{ij} . To model possible linked changes in these parameters with changes in food availability as measured by per biomass food intake rate $c_{it} = Q_{it} / B^t$ (i =juvenile index J or adult index A), we need to specify how changes in c_{it} will influence at least relative time spent foraging.

Denoting the relative time spent foraging as T_{it} measured such that the rate of effective search during any model time step can be predicted as $a_{jit} = T_{it} a_{ji}$ for each prey type j that i eats. Further, we assume that time spent vulnerable to predation, as measured by v_{ij} for all predators j on i , is inversely related to T_{it} , i.e., $v_{ij,t} = v_{ij} / T_{it}$. **An alternative structure that gives similar results is to leave the a_{ij} constant, while varying the v_{ij} by setting $v_{jit} = T_{it} \cdot v_{ij}$ in the numerator of Eq. 52 in [Predicting consumption?](#) and $v_{jit} = T_{it} \cdot v_{ij}$ in the denominator.**

For convenience in estimating the a_{ij} and v_{ij} parameters, we scale T_{it} so that $T_{i0} = 1$, and $v_{ij} = v_{ij}$. Using these scaling conventions, the key issue then becomes how to functionally relate T_{it} to food intake rate c_{it} so as to represent the hypothesis that animals with lots of food available will simply spend less time foraging, rather than increase food intake rates.

In Ecosim a simple functional form for T_{it} is implemented that will result in near constant feeding rates, but changing time at risk to predation, in situations where rate of effective search a_{ji} is the main factor limiting food consumption rather than prey behaviour as measured by v_{ji} . This is implemented in form of the relationship:

[../Resources/Images?/0800002A.png](#) Eq. 65

where, a is a user-defined Feeding time adjustment rate [0, 1] on the Ecosim [Group info?](#) form; $c_{i,opt}$ is the (internally computed) feeding rate that optimizes feeding rate versus mortality risk for i ; $c_{i,t-1}$ is the consumption/biomass ratio in the previous time step for the group. The time spent feeding is constrained by a user-defined value (Maximum relative feeding time on the *Group info* form with default of two times the feeding rate in the Ecopath base model).

The relationship between foraging time, consumption and predator biomass is illustrated in Figure 3.4.

[../Resources/Images?/0600002B.png](#)

Figure 3.4 Relationship between relative foraging time (T), Q/B and predator biomass. If Q/B is held constant the foraging time (and hence predation risk) is a linear function of the predator biomass (solid line). If T is held constant the Q/B will decrease asymptotically with predator biomass (stippled line).