

7.6 Mortality coefficients

The *Mortality coefficients form* (Figure 7.1) is one of the most important forms on the *Parameterization menu* and it is, as a rule, the first one that should be checked when balancing a model.

This form gives the Ecopath-predicted values of each type of mortality for each group in the model (see below). During balancing, the *Mortality coefficients form* will guide you in identifying where problems are (e.g., if it's fishing or predation mortality that is too high). If predation mortality is too high, then you can use the [Predation mortality](#) form to identify which predators are causing the problem.

Components of mortality in Ecopath

Under equilibrium, each group can be represented by an average organism, with an average weight. This makes it possible to use equations for estimating mortality in numbers, even when dealing with biomass. One such equation is

$$N_t = N_0 \cdot e^{-Zt} \quad \text{Eq. 25}$$

where N_0 is a number of organism at time = 0; N_t is the number of survivors at time = t ; and Z is the instantaneous rate of mortality.

Under the assumption that Z_i , the mortality of group i , is constant for the organisms included in i , it turns out that, for a large number of growth functions (including the von Bertalanffy Growth Function, or VBGF):

$$Z_i = (\text{production} / \text{biomass})_i = P/B_i \quad \text{Eq. 26}$$

or instantaneous mortality equals total production over mean biomass (Allen, 1971).

The mortality coefficient can be split into its components following a procedure well known among fisheries biologists, i.e.,

$$Z_i = P/B_i = \text{Fishing mortality} + \text{Predation mortality} + \text{Biomass accumulation} + \text{Net migration} + \text{Other mortality}$$

or

$$P/B_i = F_i + M2_i + BA_i + E_i + M0_i \quad \text{Eq. 27}$$

In some models, (e.g., the Multispecies Virtual Population Analysis model of the North Sea, Sparre, 1991), the 'other mortality' component is split between $M1_i$, i.e., predation by predators not included in the model, and $M0_i$, 'other mortality', caused by diseases, senescence, etc. In Ecopath, $M1$ is not included, as all predation mortality should be described explicitly. Further, $M0_i$ is not entered directly, but is computed from the ecotrophic efficiency, EE_i . Thus:

F_i is the Fishing mortality coefficient;

$M2_i$ is the Predation mortality coefficient;

BA_i is the Biomass accumulation coefficient;

E_i is the Net migration coefficient (immigration less emigration).

$M0_i$ is the Other mortality coefficient.

Ecopath-predicted values for these coefficients are given on the *Mortality coefficients form*. The mortality coefficients are estimated from the following equations:

$$Z_i = P/B_i$$

$$M2_i = (\sum B_j \cdot Q/B_j \cdot DC_{ji}) / B_i$$

$$F_i = Y/B_i$$

$$M0_i = (1 - EE_i) \cdot P/B_i$$

where Q/B_j is the consumption/biomass ratio of predator j ; DC_{ji} is the proportion prey i constitutes to the diet of predator j ; B_i is the average biomass of i , and C_i is the catch of i . The biomass accumulation term, BA_i , is a basic input term.

If any component of the system is harvested, a summary of the mortality coefficients can be displayed, which presents total mortality ($Z = P/B$) and its component: fishing mortality (F), other exports (E), other mortality (M_0), and predation mortality (M_2). Predation mortality is further broken down to show the contribution of each consumer groups to the total predation mortality of each prey group.

See also introductory material on [Production](#), [Consumption](#), [Dealing with open system problems](#) and [Other mortality](#).

Group name	Prod./biom (=Z)	= Fishing mort. rate	+ Predat.mort. rate	+ Biom.accum. rate	+ Net migration rate	+ Other mort. rate
Snook						
1 0-12 Snook	5.000		0.534			4.466
2 3-12 Snook	2.000		0.0159			1.984
3 12-48 Snook	0.900	0.0880				0.812
4 48-90 Snook	0.620	0.356				0.264
5 90+ Snook	0.600					0.600
Red Drum						
6 0-3 Red Drum	8.000		0.900			7.100
7 3-8 Red Drum	3.500		0.0115			3.488
8 8-18 Red Drum	1.100					1.100
9 18-36 Red Drum	0.600	0.0462				0.554
10 36+ Red Drum	0.550	0.00167				0.548
Sea Trout						
11 0-3 Sea Trout	6.000		2.199			3.801
12 3-18 Sea Trout	1.400		0.189			1.211
13 18+ Sea Trout	0.700	0.227	0.00395			0.469
Sand Trout						
14 0-3 Sand Trout	5.000		1.125			3.875
15 3-12 Sand Trout	1.200		0.514			0.686
16 12+ Sand Trout	0.700	0.165	0.000959			0.534
Mullet						
17 0-6 Mullet	6.700		0.737			5.963
18 6-18 Mullet	1.800	0.479	0.0322			1.289
19 18+ Mullet	0.800	0.179	0.000358			0.621
Mackrel						
20 Mackrel 0-3	4.000		0.385			3.615
21 Mackrel 3+	0.500	0.273				0.227
Ladyfish						
22 Ladyfish 0-10	2.800		0.318			2.482
23 Ladyfish 10+	1.600	0.112	0.0215			1.466
24 Jacks	0.600		0.360			0.240
25 Bay Anchovy	2.530	0.0519	1.466			1.012
26 Pin Fish	1.019	0.00313	0.572			0.444
27 Spot	1.100		0.180			0.920
28 Silver Perch	1.400	0.0111	1.302			0.0868

Figure 7.1 Mortality rates. Z is total mortality; F is fishing mortality; M_0 other mortality; and M_2 predation mortality. $Z = P/B = F + E + M_0 + M_2$.