

Wikiprint Book

Title: EwEugMortalityForaPreylsConsumptionForaPredator

Subject: Ecopath Developer Site - EwEugMortalityForaPreylsConsumptionForaPredator

Version: 13

Date: 2024-04-17 21:54:28

Table of Contents

2.1 Mortality for a prey is consumption for a predator	3
--	---

2.1 Mortality for a prey is consumption for a predator

The core routine of Ecopath is derived from the Ecopath program of Polovina (1984a; 1984b) modified to render superfluous its original assumption of steady state. Ecopath no longer assumes steady state but instead bases the parameterization on an assumption of mass balance over an arbitrary period, usually a year (but also see discussion about seasonal modelling). In its present implementation Ecopath parameterizes models based on two master equations, one to describe the production term and one for the energy balance of each group.

The first Ecopath equation describes how the production term for each group (i) can be split in components. This is implemented with the equation,

Production = catches + predation mortality + biomass accumulation + net migration + other mortality **Eq. 1**

or, more formally,

$$P_i = Y_i + B_i M2_i + E_i + BA_i + P_i(1 - EE_i) \text{ Eq. 2}$$

where P_i is the total production rate of (i), Y_i is the total fishery catch rate of (i), $M2_i$ is the total predation rate for group (i), B_i the biomass of the group, E_i the net migration rate (emigration - immigration), BA_i is the biomass accumulation rate for (i), while $MO_i = P_i \cdot (1 - EE_i)$ is the 'other mortality' rate for (i).

This formulation incorporates most of the production (or mortality) components in common use, perhaps with the exception of gonadal products. Gonadal products however nearly always end up being eaten by other groups, and can be included in either predation or other mortality.

Eq. 2.2 can be re-expressed as

,

$$P/B_i = Q/B_i + DC_{ji} \text{ Eq. 3}$$

$$P/B_i = Q/B_i + DC_{ji} \text{ Eq. 4}$$

where: P/B_i is the production/biomass ratio, Q/B_i is the consumption/biomass ratio, and DC_{ji} is the fraction of prey (i) in the average diet of predator (j).

Based on Eq. 2.3, for a system with n groups, n linear equations can be given, in explicit terms,

$$P/B_i = Q/B_i + DC_{ji} \text{ Eq. 5}$$

This system of simultaneous linear equations can be re-expressed

$$P/B_i = Q/B_i + DC_{ji} \text{ Eq. 6}$$

with n being equal to the number of equations, and m to the number of unknowns.

This can be written in matrix notation as

$$[A]_{nm} \cdot [X]_m = [Q]_m \text{ Eq. 7}$$

Given the inverse A^{-1} of the matrix A , this provides

$$[X]_m \cdot [A^{-1}]_{n,m} = [Q]_m \text{ Eq. 8}$$

If the determinant of a matrix is zero, or if the matrix is not square, it has no ordinary inverse. However, a generalized inverse can be found in most cases (Mackay, 1981). In the Ecopath model, the approach of Mackay (1981) is used to estimate the generalized inverse.

If the set of equations is over-determined (more equations than unknowns), and the equations are not consistent with each other, the generalized inverse method provides least squares estimates, which minimizes the discrepancies. If, on the other hand, the system is underdetermined (more unknowns than equations), an answer that is consistent with the data will still be output. However, it will not be a unique answer.

Of the terms in Eq. 2.3, the production rate, P_i is calculated as the product of B_i the biomass of (i) and P_i/B_i the production/biomass ratio for group (i). The P_i/B_i rate under most conditions corresponds to the total mortality rate, Z , see Allen (1971), commonly estimated as part of fishery stock assessments. The 'other mortality' is a catch-all term including all mortality not elsewhere included, e.g., mortality due to diseases or old age, and is internally computed from,

$$MO_i = P_i \cdot (1 - EE_i) \text{ Eq. 9}$$

where EE_i is called the 'ecotrophic efficiency' of (i), and can be described as the proportion of the production that is utilized in the system. The production term describing predation mortality, $M2$, serves to link predators and prey as,

[../Resources/Images?/0800000B.png](#) Eq. 10

where the summation is over all (n) predator groups (j) feeding on group (i), Q_j is the total consumption rate for group (j), and DC_{ji} is the fraction of predator (j)'s diet contributed by prey (i). Q_j is calculated as the product of B_j , the biomass of group (j) and Q_j/B_j , the consumption/biomass ratio for group (j).

An important implication of the equation above is that information about predator consumption rates and diets concerning a given prey can be used to estimate the predation mortality term for the group, or, alternatively, that if the predation mortality for a given prey is known the equation can be used to estimate the consumption rates for one or more predators instead.

For parameterization Ecopath sets up a system with (at least in principle) as many linear equations as there are groups in a system, and it solves the set for one of the following parameters for each group:

Biomass; Production/biomass ratio; Consumption/biomass ratio; or Ecotrophic efficiency.

If, and only if, all four of these parameters are entered, the program will prompt you during basic parameterization whether to estimate the biomass accumulation, or, alternatively, to estimate the net migration rate. If a positive response is given, the program will use all the four basic parameters and it will establish mass-balance by calculating one of the two other parameters. If only three of the basic parameters are entered the following parameters must be entered for all groups:

Catch rate; Net migration rate; Biomass accumulation rate; Assimilation rate; and Diet compositions.

It was indicated above that Ecopath does not rely on solving a full set of linear equations, i.e., there may be less equations than there are groups in the system. This is due to a number of algorithms included in the parameterization routine that will try to estimate iteratively as many 'missing' parameters as possible before setting up the set of linear equations. The following loop is carried out until no additional parameters can be estimated.

The net growth efficiency, g_p , is estimated using

$$cg = (P/B_i) / (Q/B_i) \quad \text{Eq. 11}$$

while P/B_i and Q/B_i are attempted solved by inverting the same equation. The P/B ratio is then estimated (if possible) from

[../Resources/Images?/0800000C.png](#) Eq. 12

This expression can be solved if both the catch, biomass and ecotrophic efficiency of group i , and the biomasses and consumption rates of all predators on group i are known (including group i if a zero order cycle, i.e., 'cannibalism' exists). The catch, net migration and biomass accumulation rates are required input, and hence always known;

The EE is estimated from

[../Resources/Images?/0800000D.png](#) Eq. 13

where the predation mortality $M2$ is estimated from Eq. 2.2.10.

In cases where all input parameters have been estimated for all prey for a given predator group it is possible to estimate both the biomass and consumption/biomass ratio for such a predator. The details of this are described in Appendix 4, Algorithm 3.

If for a group the total predation can be estimated it is possible to calculate the biomass for the group as described in detail in Appendix 4, Algorithm 4.

In cases where for a given predator j the P/B , B , and EE are known for all prey, and where all predation on these prey apart from that caused by predator j is known the B or Q/B for the predator may be estimated directly.

In cases where for a given prey the P/B , B , EE are known and where the only unknown predation is due to one predator whose B or Q/B is unknown, it may be possible to estimate the B or Q/B of the prey in question.

After the loop no longer results in estimate of any 'missing' parameters a set of linear equations is set up including the groups for which parameters are still 'missing'. The set of linear equations is then solved using a generalized inverse method for matrix inversion described by Mackay (1981). It is usually possible to estimate P/B and EE values for groups without resorting to including such groups in the set of linear equations.

The loop above serves to minimize the computations associated with establishing mass-balance in Ecopath. The desired situation is, however, that the biomasses, production/biomass and consumption/biomass ratios are entered for all groups and that only the ecotrophic efficiency is estimated, given that no procedure exists for its field estimation.